

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
24 October 2002 (24.10.2002)


PCT

(10) International Publication Number
WO 02/083030 A2

- (51) International Patent Classification⁷: **A61F**
- (21) International Application Number: **PCT/US02/11476**
- (22) International Filing Date: **12 April 2002 (12.04.2002)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/284,308 17 April 2001 (17.04.2001) US
10/066,323 31 January 2002 (31.01.2002) US
- (71) Applicant: **CRYOLIFE, INC.** [US/US]; 1655 Roberts Boulevard, N.W., Kennesaw, GA 30144 (US).
- (72) Inventors: **GILBERT, Carl, W.**; 4655 Oakleigh Manor Drive, Powder Springs, GA 30127 (US). **MCGOWAN, Eleanor, B.**; 3062-I Spring Hill Parkway, Smyrna, GA 30080-4735 (US). **BLACK, Kirby, S.**; 1371 Peppergrass Trail, Acworth, GA 30101 (US). **HARPER, T., Gregory, P.**; 86 West Euclid Avenue, Stockton, CA 95204 (US).
- (74) Agent: **MERCANTI, Michael, N.**; Roberts & Mercanti, L.L.P., P.O. Box 3156, 105 Lock Street, Suite 203, Newark, NJ 07103-0156 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report

[Continued on next page]

(54) Title: BIFUNCTIONAL ENERGY-REVERSIBLE ACYL-COMPOSITIONS



(57) Abstract: Energy-reversible acyl conjugates, intermediates, and related compositions are disclosed. In preferred aspects, examples of such compositions include Formulae (A) and (B).

WO 02/083030 A2



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

BIFUNCTIONAL ENERGY-REVERSIBLE ACYL-COMPOSITIONS

Background of the Invention

The present application relates generally to energy reversible acyl-enzymes. In particular, the present application relates to energy reversible acyl-enzymes and the like having a cinnamate or related structure core and an additional reactive group which can be modified to impart new properties to the whole composition.

Various methods to make use of enzyme inactivation have been disclosed. Enzyme inhibition can be used to enhance long term storage of enzymes or to inactivate the enzymes in a pharmaceutical drug. For example, U.S. Patent No. 5,770,699 describes a process of enzyme inhibition to produce inactivated blood factors. U.S. Patent No. 5,837,679 discloses a method to extend the half lives of blood factors via a transient modification of blood factors by acylation. U.S. Patent No. 4,337,244 reports a method of treating venous thrombosis using an inactivated fibrinolytic enzyme.

Enzyme activity can be controlled with inhibitors. Reversible control of enzyme activity with light has been the focus of a number of reports (see U.S. Patent Nos. 5,114,851 and 5,218,137 to Porter et al.). There are a number of advantages of this concept. Most striking is the ability to control enzyme activity specifically and rapidly, by exposure to light *in vivo* or *ex vivo*.

Porter et al. disclose in U.S. Patent Nos. 5,114,851 and 5,218,137, the light controllable enzymes are obtained by coupling an enzyme active site amino acid residue to cinnamate (CINN) derivatives to form *o*-hydroxy cinnamate substituted esters or acyl enzymes, which are inactive. On photolysis, the bond with the active site amino acid residue is cleaved and the active site is exposed. Pizzo et al. [(1986) Ann. N.Y. Acad. Sci. 485: 199-203] reported on the use of an *o*-hydroxy cinnamate substituted ester, formed by coupling the active site of an enzyme to α -methyl-2-hydroxy-4-diethylaminocinnamic acid.

A subsequent report by Porter, et al. demonstrated the therapeutic potential of the inhibited enzymes. Control of clotting reaction times was concentration dependent and photolysis time dependent. *In vivo* clotting of abnormal blood vessels in a rabbit model of corneal neovascularization was achieved by injection of the inhibited, "caged" enzymes

and application of 366 nm light to the eyes for 25 minutes. See Arroyo et al., *Thromb. Haemost.* 78, 791-793 (1997). U.S. Patent Nos. 5,114,851 and 5,218,137, further describe these cinnamate derivatized enzymes and uses thereof.

5 The usefulness of the inhibited enzymes is a function of the rate and extent of photolysis. Utility is limited if rate of bond cleavage between the cinnamate moiety and the active-site amino acid is slow. Many applications require rapid exposure times, on the order of seconds in most *in vitro* or typical applications, or at most minutes, in most *in vivo* applications. Rapid, controlled response times are essential for most clinical applications and are of particular importance with labile enzymes or uses where rapid
10 reaction times are essential, as in clotting.

For example, the formation of an acyl-enzyme between α -chymotrypsin and the *p*-nitrophenyl ester of *p*-nitro-*trans*-cinnamic acid is described by Varfolomeyev, S., et al., [*FEBS Lett.* 15:118 (1971)]. The bond between the enzyme and the carboxylate group is formed with the hydroxyl group of the serine at the catalytic center of the enzyme
15 [Berezin, I. et al., *FEBS Lett.* 8:173 (1970)]. The formation of an acyl-enzyme between α -thrombin and the *trans*-isomer of an ester of *o*-hydroxy- α -methylcinnamic acid is described by Turner, A., et al., *J. Amer. Chem. Soc.* 109:1274 (1987). The bond between the enzyme and the carboxylate group is formed with the hydroxyl group of the serine-195 at the catalytic center of the enzyme (Turner, A., et al., *J. Amer. Chem. Soc.* 110:244
20 (1988)). Exposure of the compound to light led to deacylation. The photoactivation of these enzymes was slow and required light intensities and wavelengths such that appreciable enzyme degradation occurred during photoactivation.

Research in this area has continued. For example, Porter, et al. (*Photochem. Photobiol. B* 38(1), 61-69 (1997) inserted a biotin derivative on the 2-position of the
25 cinnamate side chain (adjacent to the carboxylate group), which could be bound to avidin, for purposes of purification and immobilization. The modified compound also maintained the ability to be photoactivatable.

Further modifications of the CINN core molecule would be desirable in order to improve inhibited enzyme compositions that can be rapidly and controllably reactivated.
30 For example, it would be desirable to introduce additional sites to react with other molecules for purposes of changing the properties of the acyl-enzyme (such as immobilization or pharmacokinetics), while maintaining the desired properties of photoactivation. The present invention addresses this need.


Summary of the Invention

It is an object of the present invention to provide improved compositions designated herein as Z-CINN-X₁-A, including inhibited enzymes that can be rapidly and controllably reactivated.

5 It is another object of the present invention to provide an additional site on Z-CINN-X₁-A type compounds (as defined below) which can be derivatized in various ways, including with groups designated herein as "B-L", to provide compositions designated herein as B-L-Z-CINN-X₁-A, with additional properties such as stability, targeting capacity, or immobilization to appropriate supports.

10 It is yet another object of the present invention to provide an enzyme inhibitor composition (B-L-Z-CINN-X₁-A₁), which is capable of generating an inactivated enzyme composition (B-L-Z-CINN-X₁-A₂), where the enzyme can be released in active form via a controlled mechanism such as input of light energy.

15 These and other objects are provided by the present invention which in one embodiment provides compositions corresponding to Z-CINN-X₁-A and Formula (I):




wherein:

R₁ and R₂ are individually selected from among H, CH₃, C₂-C₁₀ alkyls, alkenyls or alkynyls which can be substituted or unsubstituted; straight or branched; C₂-C₁₀ heteroalkyls, heteroalkenyls or heteroalkynyls and -(CR₁₅R₁₆)_p-D,

wherein: R₁₅ and R₁₆ are individually selected from the group consisting of H, CH₃, C₂-C₁₀ alkyls, alkenyls or alkynyls which can be substituted or unsubstituted; straight or branched; C₂-C₁₀ heteroalkyls, heteroalkenyls or heteroalkynyls;

25 p is a positive integer from 1 to about 12;

D is selected from among -SH, -OH, X₂, -CN, -OR₁₉, NHR₂₀,



wherein:

R₁₇ is H, CH₃ or X₃;

R₁₈ is H, a C₁-C₄ alkyl or benzyl;

5 R₁₉ is H, a C₁₋₄ alkyl, X₂ or benzyl;

R₂₀ is H, a C₁₋₁₀ alkyl or -C(O)R₂₁,

wherein R₂₁ is H, a C₁₋₄ alkyl or alkoxy, t-butoxy or benzyloxy;

X₂ and X₃ are independently selected halogens;


10 R₃ is H, CH₃, or -C(=O)(CR₁₅R₁₆)_w-D,

where w is 0 or an integer from 1 to about 12, and D is H or as described for R₁ and R₂

J is O, NH or S;

15 R₄, R₅, and R₆ are independently selected from among H, CH₃, C₂-C₁₀ alkyls, alkenyls or alkynyls which can be substituted or unsubstituted; straight or branched; C₂-C₁₀ heteroalkyls, heteroalkenyls or heteroalkynyls and halogens;

Z is NR₇R₈ or



wherein R₇ is selected from among H, CH₃, C₂-C₁₀ alkyls, alkenyls or alkynyls which can be substituted or unsubstituted; straight or branched; C₂-C₁₀ heteroalkyls,

20 heteroalkenyls or heteroalkynyls, or -(CR₂₃R₂₄)_q-aryl, or R₈;

wherein R₂₃ and R₂₄ are independently H or a C₁-C₁₀ alkyl;

q is an integer from 1 to about 6;

R₈ is selected from among (CR₉R₁₀)_n-NR₂₂-R₁₁, (CR₉R₁₀)_n-CH₂-NHC(O)R₂₆ and (CR₉R₁₀)_n-CH₂-E,

25 wherein R₉ and R₁₀ are independently selected from among H, CH₃, C₂-C₁₀ alkyls, alkenyls or alkynyls which can be substituted or unsubstituted; straight or branched;

C_2-C_{10} heteroalkyls, heteroalkenyls or heteroalkynyls and halogens;

R_{26} is H, CH_3 , O-t-butyl, O-benzyl;

E is OH, SH or $O-C(O)R_{27}$,

wherein R_{27} is a C_1-C_6 alkyl, benzyl or phenyl;

5 R_{22} is H or CH_3 ;

n is a positive integer from 1 to about 10;

R_{11} is H or -L-B

wherein L is a linker; and

B is a first active moiety, containing reactive group moieties such

10 as maleimidyl or N-hydroxysuccinimidyl, etc., or a polymer;

R_{25} is H, $-C(O)-R_{28}$ or $-C(O)-O-R_{29}$,

wherein R_{28} is a C_1-C_6 alkyl or benzyl; and R_{29} is CH_3 , t-butyl or benzyl;

X_1 is O, NH, or S; and

A is H or a second active moiety, designated as either A_1 or A_2 herein, it being


15 understood that X_1 is an integral part of either A_1 or A_2 .

Pharmaceutically acceptable salts, including Cl^- , Br^- , HSO_4^- , etc. of compounds corresponding to Formula (I) are also provided.

In preferred aspects, B is a polymer or natural or synthetic organic molecule, a protein, such as an antibody or fragment thereof, a carbohydrate, a nucleic acid, a lipid, 20 or other naturally occurring or synthetic compounds.

In further preferred embodiments, A is A_1 , and X_1A_1 is a substrate or substrate analog such as an amino acid, an amino acid derivative, a peptide, a peptide derivative or a substrate or substrate analog for serine proteases, cysteine proteases, esterases, lipases, or other enzymes containing an active site serine or cysteine. In some preferred aspects,

25 X_1A_1 is:



in which R_{12} and R_{13} are H or electron donating or electron withdrawing groups and W is a covalent bond. Alternatively, A_1 can also be an amino acid or amino acid derivative, peptide or peptide derivative or other molecule which is a substrate or substrate analog for 30 serine proteases, cysteine proteases, esterases, lipases, or other enzymes containing an active site serine or cysteine.

In other aspects of the invention, $H-X_1A_2$ is an enzyme such as a serine protease, cysteine protease, esterase, lipase, or other enzyme containing an active-site serine, cysteine or lysine whose side-chain $-O$, $-S$ or $-NH$ corresponds to X_1 of Formula (I) which is bonded to the $C(O)$ of the Z-CINN upon displacement of X_1A_1 . Thus, $H-X_1A_2$ is an enzyme rendered inactive through its bond to Z-CINN and is preferably capable of having its enzyme activity restored by hydrolysis or by exposure to light or other energy source.

In still further aspects of the invention, Z-CINN- X_1 - A_1 compositions are derivatized utilizing the reactive Z site of Formula I. In particular, when R_8 is $(CR_9R_{10})-NR_{22}-R_{11}$, and R_{11} is L-B, the artisan is provided with Z-CINN cores which are linked to, among other things, polymers such as PEG or other activated polymers.

Alternatively, the Z-CINN- X_1 -A can be joined to a linker group L so that subsequent attachment of B groups, e.g. monoclonal antibodies (mAbs), polymers, etc. can be carried out when desired. The Z-CINN- X_1 -A can be linked using, for example, succinimidyl-6-[(β -maleimidopropionamido)hexanoate], ethylene glycol bis[succinimidyl succinate], bis activated PEGs containing terminal succinimidyl succinates, N-hydroxy-succinimidyls and/or maleimides. Other bifunctional reagents are also contemplated. The L-Z-CINN- X_1 -A molecules can then be reacted with another molecule, e.g. proteins, including antibodies, fragments thereof, nucleic acids, lipids, or other naturally occurring or synthetic compounds. Derivatized compositions (B-L-Z-CINN- X_1 - A_1) can then be reacted with an enzyme to form B-L-Z-CINN- $X-A_2$, where the enzyme is inactivated.

As a result of the present invention, several advantages are provided. For example, these Z-CINN- X_1 -A inactivated compositions can have various beneficial properties, such as increased solubility, increased half-life in circulation, targeting ability, or other features. In addition, L-Z-CINN- X_1 -A inactivated compositions can be immobilized by crosslinking to support materials via the linker L. The support materials can be any industrially or pharmaceutically suitable materials such as organic polymers, inorganic polymers, natural polymers, biopolymers or zeolites and can be in the form of films, membranes, filters, beads, particles, resins, microparticles, or columns. Alternatively, B-L-Z-CINN- X_1 -A can be attached to supports by mechanisms such as affinity or by additional coupling reactions with activated support materials.

The acyl-enzyme bonds can be relatively stable or susceptible to hydrolysis at about neutral pH in the dark. They are susceptible to cleavage by a source of energy such as light, including ultraviolet, visible, and infrared light, microwave, ultrasound, radiowave energy or radioactivity. The preferred energy source is light having a

wavelength in the range from 340 to 700 nm and preferably 350 to 420 nm. This provides, for example, a means to controllably release and convert inactivated, acylated enzymes to active enzymes.

Compounds of the invention corresponding to, for example, B-L-Z-CINN-X₁-A₂, can be used in an assay, where the acyl-enzyme is first free or bound to an immobilized support, then energy is applied to release the active enzyme into solution. The inactivated compositions can also be used in purification methods, and as therapeutics, which are activated at the time of or shortly before administration. The inactivated enzymes can be stored for lengthy periods, then reactivated at time of use, to increase shelf lives.

In further aspects of the invention, methods of preparing and using the compositions of the present invention are also provided.

Brief Description of the Drawings

Figure 1 illustrates the synthesis of (3-[2-hydroxy-4-(ethyl)(2-tert-butylloxycarbamidoethyl)amino]phenyl-2-methyl-2-propenoic acid, ethyl ester) (10).

Figure 2 illustrates the synthesis of (3-[2-hydroxy-4-(ethyl)aminoethyl)-amino]phenyl-2-methyl-2-propenoic acid, 4-aminoiminophenyl ester, HCl salt) (N-CINN-AP) (16).

Figure 3 illustrates the preparation of mPEG₅₀₀₀-N-CINN-AP (18).

Figure 4 illustrates the preparation of derivatized inhibited enzyme (19) by reaction of mPEG₅₀₀₀-N-CINN-AP (18) and enzyme.

Figure 5 illustrates the preparation of a maleimido derivative of N-CINN-AP (20), Mal-L-N-CINN-AP, by reaction of (16) with succinimidyl-6-[(β-maleimidopropion-amido)hexanoate].

Figure 6 illustrates the preparation of antibody-L-N-CINN-AP (21) by reaction of SH-modified protein with (20), followed by reaction with enzyme to give antibody-Mal-L-N-CINN-enzyme (22).

Detailed Description of the Invention

I. Definitions

The term "inhibitor" as used herein means a compound which can react with a residue at the catalytically active site of an enzyme to inactivate the enzyme by binding to the site and preventing catalytic activity.

The term "inactivated" as used herein means that the catalytically active site of an enzyme is covalently coupled to an inhibitor.

The term "reactivation" as used herein refers to the process in which the covalent acyl bond is cleaved by hydrolysis, or by a source of energy, and enzyme activity is restored.

The term "cinnamic acid" is the common or trivial name for 3-phenyl-2-propenoic acid. **5** acid. Z-CINN-X₁-A molecules include derivatives of cinnamic acid.

IUPAC names are given for synthesized compounds.

For purposes of the present invention "reactive group", particularly as used to describe the variable B shall be understood to include those moieties capable of facilitating conjugation of L and one or more biologically active moieties or polymers, including but **10** not limited to the free electron pair (double bond) of a maleimidyl residue.

For purposes of the present invention, the terms "alkyl", "alkenyl" and "alkynyl" shall be understood to include, e.g., straight, branched, substituted, cyclo- derivatives thereof, including alkoxy or substituted cycloalkyls, etc. "Substituted" shall be understood to include halo-, alkoxy-, nitro-, etc.

15 For purposes of the present invention, "molecule" and "biologically active molecule" shall be understood to embrace not only organic or small molecules but also proteins, peptides and the like.


For purposes of the present invention, "halogen" shall be understood to include chlorine, fluorine, bromine, etc.

20

II. Z-CINN Derivatives

A. Z-CINN Various compositions can be formed using Z-CINN as a core molecule. A variety of functional groups can be introduced into Z-CINN to modify its chemical as well as physical properties. These compositions can also have different **25** reactivities toward a nucleophile or light. Some preferred inhibited compositions are discussed in detail as follows.

The Z-CINN core molecule corresponds to Formula (II)




wherein all variables are as set forth in Formula (I) above.

30

In one preferred embodiment for Formulas (I) and (II), Z is NR_7R_8 . Further, R_1 , R_2 , R_3 , R_4 , R_5 , and R_6 are H or lower alkyl groups such as CH_3 or CH_2CH_3 . In another preferred embodiment for Formulas (I) and (II), R_7 is CH_3CH_2 and R_8 is $-(\text{CR}_9\text{R}_{10})_n-\text{NR}_{22}-\text{R}_{11}$ and R_9 and R_{10} are H; wherein n is 2; R_{22} is H or CH_3 ; and X_1 is O, S or N. In those aspects of the invention where J is O and Z is NR_7R_8 , some preferred R_8 groups include:

1. $(\text{CH}_2)_n-\text{NH}_2$ and salts thereof, e.g. $-\text{NH}_3^+:\text{Cl}^-$ or Br^- or HSO_4^- , etc.
2. $(\text{CH}_2)_n-\text{NH}-\text{C}(\text{O})-\text{H}$
3. $(\text{CH}_2)_n-\text{NH}-\text{C}(\text{O})-\text{CH}_3$
- 10 4. $(\text{CH}_2)_n-\text{NH}-\text{C}(\text{O})-\text{O}-t\text{-butyl}$
5. $(\text{CH}_2)_n-\text{NH}-\text{C}(\text{O})-\text{O}-\text{benzyl}$
6. $(\text{CH}_2)_n-\text{OH}$
7. $(\text{CH}_2)_n-\text{SH}$
8. $(\text{CH}_2)_n-\text{O}-\text{C}(\text{O})-\text{R}_{21}$; wherein R_{21} is CH_3 , C_{1-6} alkyl or phenyl;
- 15 wherein each of the above n is an integer of from 2 to about 10, preferably 2-4.

In still a further aspect of the invention, Z is substituted piperazine thereby providing compositions of the Formula (Ia):



20

wherein all the variables are as previously defined, with regard to Formula (I).

The cinnamic acid backbone enzyme inhibitors can be synthesized using standard techniques and available reagents by those skilled in the art. For example, as Figure 1 demonstrates, an amine such as 2-ethylaminoethylamine (1) can react with an anhydride (2) to generate amide (3). 3 condenses with 1,3-cyclohexanedione (4) followed by dehydration to generate 3-amino-2-cyclohexen-1-one (5). Dehydrogenation of 5 in the presence of a catalyst generates a 3-hydroxyphenyl amine (6). Hydrolysis removes a blocking group to obtain 7, followed by N-acylation to obtain 8. Reaction of 8 with POCl_3/DMF generates an aldehyde (9). Reaction of 9 with an appropriate Wittig reagent, carbethoxyethylidene triphenylphosphorane (10), generates 11, where the aldehyde group

30

is converted to a 2-propenoic acid ethyl ester. The N-acyl group is exchanged to obtain 12, which is hydrolyzed in LiOH to obtain 13 (tBOC-N-CINN). The compounds of Formula (Ia) can be similarly prepared using piperazine in place of the amine (1).

As Figure 2 further demonstrates, 13 can be reacted in the presence of an alcohol such as 4-aminoiminophenol (14), dicyclohexylcarbodiimide, and dimethylaminopyridine to form an ester (15). The N-protective group can be hydrolyzed in the presence of HCl to generate a substituted cinnamate ester (enzyme inhibitor) containing a free, protonated amino group (16). As Figure 3 demonstrates, 16 further reacts with a monomeric SS-PEG (17) to generate PEG-derivatized inhibitor (18).

Figure 4 further illustrates the reaction of 18 with various enzymes containing active site serines and affinity for 4-aminoiminophenol to obtain inhibited enzyme compositions.

In another embodiment, other alcohols, thiols, or amino compounds, such as amino acids, amino acid derivatives, peptides, or other compounds are used in place of 14 to form additional $-X_1A_1$ moieties. For example, an amino acid derivative or peptide derivative is coupled to the N-CINN backbone (13) via the DCI/DMAP chemistry shown in Figure 2. Carboxyl groups on a peptide or amino acid would be protected with acceptable blocking groups prior to reaction with 13. The amino group would then react with 13, DCI, and DMAP to give an amide bond, and the blocking groups would be removed. Similar procedures would be followed for alcohols or thiols.

Figure 5 is the general scheme for preparation of a maleimido derivative of N-CINN-AP (20), by reaction of (16) with a heterobifunctional reagent containing N-hydroxysuccinimide and maleimide. An amino group of 16 reacts with a reagent such as SMPH (succinimidyl-6-[β -maleimidopropionamido] hexanoate; Pierce); the maleimide group at the other end is available for further reactions. A number of similar reagents are available, for example bifunctional NHS, bifunctional PEGs, or similar compounds. Those skilled in the art will be able to choose appropriate linking reagents to couple the Z-CINN backbone to carrier groups without undue experimentation. For purposes of illustration and not limitation, a non-limiting list of linkers which can be employed include succinimides, maleimides, imidoesters, 2-iminothiolane, hydrazides, maleic anhydride, azides, citraconic anhydride, glutaraldehyde, and the like, which facilitate attachment of any desired B group to the Z-CINN core.

Figure 6 shows that Mal-L-N-CINN-AP (20) can be reacted with a SH-modified antibody to form an antibody-L-N-CINN-AP (21). 21 can be reacted with free enzyme to

form antibody-L-N-CINN-enzyme (22), where the enzyme is inhibited. 22 can be treated with light or other energy source to regenerate free enzyme. SH-modified antibody is prepared at ~pH 7 using common reagents and the Mal-L-N-CINN-AP is added at ~pH 7 in slight excess. Free reagents are separated by ultrafiltration or size exclusion chromatography. In other embodiments, moieties other than SH-modified antibodies are used following the schematic set forth in Figure 6, with changes in the reactive linking group as required.

B. Inhibition of Enzyme Activity

As defined by Formula I, X_1A may include an active moiety. In that formula, the active moiety is referred to as the "second" so as to distinguish it from the first active moiety which is part of B and is discussed in more detail below. Thus, X_1A_2 can be a biologically active moiety such as an enzyme which has a serine or cysteine residue at the catalytically active site, such as a number of proteases or esterases. The enzyme $H-X_1A_2$ can be specifically reacted with Z-CINN- X_1-A_1 , a substrate or substrate analog for the enzyme. A transition state forms, where the enzyme displaces the leaving group in the acyl bond, and an inhibited acyl-enzyme is formed between the Z-CINN and the enzyme, and is designated herein as Z-CINN X_1-A_2 . The inhibited enzyme can be reactivated and recovers its activity when the acyl bond is cleaved by hydrolysis or photolysis.


Exemplary enzymes ($H-X_1-A_2$) are serine proteinases (proteases) such as chymotrypsin, trypsin, acrosin, clotting factors such as thrombin, VIIa, IXa, Xa, XIa or XIIa, cathepsin G, fibrinolytic pathway proteins such as plasmin, tissue plasminogen activator, urokinase, streptokinase, and complement proteins such as C3/C5 convertase, complement Factor I, and complement Factor D.

Other exemplary enzymes include cysteine proteinases such as cathepsin B, papain, and bromelain, as well as lipases, and esterases such as acetylcholinesterase.

Preferred serine proteinases include chymotrypsin, trypsin, thrombin, plasmin, acrosin, coagulation factors IXa, Xa, XIa, and XIIa, plasminogen activator, plasma kallikrein, tissue kallikrein, urokinase, plasmin, pancreatic elastase, and leukocyte elastase. Inhibitors are preferably target specific. For example, imidazole derivatives are known as good substrates for α -chymotrypsin. F. Markwardt et al., *Pharmazie* 29:333 (1974); G. Wagner et al., *Pharmazie* 28:293 (1973); J. Sturzebecher et al., *Acta Biol. Med. Germ.* 35:1665 (1976); P. Walsmann, *Folia Haematol., Leipzig* 109:75 (1982); V. Valenty et al., *Biochem. Biophys. Res. Comm.* 88:1375 (1979); and F. Markwardt et al., *Acta Biol. Med.*

- Germ* 28:19 (1972)) describe compounds that lead to stable carboxylate esters of the enzyme active site serine. Other inhibitors that have been studied include compounds that react with the enzyme to generate stable sulfonate or phosphate esters (See R. Laura et al., *Biochemistry* 19:4859 (1980); S. Wong and E. Shaw, *Arch. Biochem. Biophys.* 161:536 (1974)).
- 5 Peptide derivatives of phosphonates have also been used (J. Oleksyszyn and JC Powers, *Biochemistry* 15: 485 (1991)).

Alternatively, A can be a moiety designated herein as A₁. Preferred X₁A₁ groups include




- 10 in which R₁₂ and R₁₃ are electron donating or electron withdrawing groups and W is N or CH, or X₁A₁ is an amino acid or amino acid derivative, peptide or peptide derivative, or other substrate or substrate analog of enzymes such as serine proteases, cysteine proteases, or esterases containing an active-site serine or cysteine that can be cleaved by the enzyme, leading to binding of the enzyme to the cinnamoyl moiety as an acyl-enzyme.
- 15 Z-CINN-X₁A₁ represent inhibitors of the enzymes discussed above. Of the preferred inhibitors, those in which *p*-nitrophenyl is the leaving group (X₁A₁) are preferred for use in making acyl-enzymes with enzymes such as chymotrypsin. Those inhibitors in which either 4-aminoiminophenyl or 4-guanidinophenyl are the leaving groups are preferred for use in making acyl-enzymes with coagulation enzymes, such as thrombin,
- 20 plasmin, coagulation Factor IXa, Xa, XIa, XIIa, or with tissue plasminogen activator, urokinase, and trypsin. Those in which amino acids or amino acid derivatives, peptides or peptide derivatives, or other substrates or substrate analogs are the leaving groups will react with a variety of enzymes containing serine or cysteine at the active site, including, for example, proteases, lipase, esterases, etc.

25

C. Derivatives of Z-CINN Backbones

A number of different molecules can be attached to the Z-CINN backbone of the present invention. See Formula (III) below.



wherein all variables are as previously defined.

In one aspect, moieties can be attached to increase solubility or circulating half-life. For example, an activated polyethylene glycol (PEG) or polypropylene glycol (PPG) can be attached to the Z-CINN core molecule. See U.S. Patent No. 4,179,337 generally and U.S. Patent No. 5,612,460 which describes *inter alia* poly(ethylene glycol)-N-succinimidyl carbonate and derivatives thereof. See also, Poly(ethylene glycol) Chemistry and Biological Applications, Harris, et al., 1997 ACS. The contents of each of the foregoing is incorporated herein by reference. Generally, with reference to Figure 3, an activated mPEG (for example SS-mPEG₅₀₀₀, Shearwater Polymers, Inc.) is reacted with 16 at ~pH 7.0 for 60 min. Excess SS-mPEG is quenched with glycylglycine and the PEG-N-CINN-AP (18) is reacted with an active serine protease at pH 6.5-7.0 for about 12-16 hours at ambient temperature to give PEG-N-CINN-acyl-enzyme (19). This complex can be included as part of a pharmaceutically acceptable solution and administered to a patient in need of such therapy. The conjugate is activated with light to release native enzyme when needed. It will be appreciated by those of ordinary skill that the specific type of activated PEG or other polymer employed will be dependent upon the particular needs of the artisan and the final product desired. It is contemplated that most commercially available activated polymers are useable herein without undue experimentation.

More specifically, in order to form the polymer conjugates of the invention, R₁₁ may comprise a polymer. For example, polymers such as polyalkylene oxides (PAOs) or similar biologically acceptable polymers are converted into activated forms, as that term is known to those of ordinary skill in the art. Thus, one or both of the terminal polymer hydroxyl end-groups, (i.e. the alpha and omega terminal hydroxyl groups) are converted into the same (homo-) or different (hetero-) reactive functional groups that allow covalent conjugation to the Z-CINN as part of R₁₁, and, if desired another active or targeting group.

Other substantially non-antigenic polymers are similarly "activated" or functionalized. As an alternative to PAO-based polymers, other substantially non-antigenic or effectively non-antigenic materials such as collagen, glycosaminoglycans, poly-aspartic acid, poly-L-lysine, poly-lactic acid, copolymers of the foregoing including polylactic-polyglycolic acid
 5 copolymers, poly-N-vinylpyrrolidone, collagen cross-linked to hydrophilic polymers or any other suitable non-reactive polymer such as polyethylene alcohols can be used. Specifically preferred polymeric groups are mono or bifunctionally activated polyethylene glycol (PEG) based polymers.

Polymeric groups of any molecular weight range are usable. A preferred polymer
 10 molecular weight can range from 2,000 Daltons to 200,000 Daltons. A more preferred molecular weight range is between 5,000 to 50,000 D. The most preferred polymer molecular weight range is between 12,000 and 40,000 D, (number average).

In another embodiment, a first active molecule (B) is attached to L-Z-CINN- X_1A_1 . For example, a protein such as a monoclonal antibody (mAb) can be
 15 linked to a Z-CINN- X_1A_1 . For example, 16 is reacted with a bifunctional reagent containing both NHS and Mal (Pierce Chemical Co.) at pH ~7.0 for 1-hour to give 20 (Mal-L-N-CINN-AP). Excess NHS-Mal reagent is quenched with glycylglycine. 20 is then reacted with SH-modified antibody at pH 7.0 for 2 hours at ambient temperature to give 21 (mAb-L-N-CINN-AP). 21 is freed of excess reagent 20 by ultrafiltration or gel
 20 filtration chromatography and can then be reacted with a serine protease at pH 6.5 for about 12 to 16 hours to give 22 (mAb-L-N-CINN-acyl-enzyme). 22 (B-L-N-CINN- X_1A_2) can be targeted to a site with the mAb, then active serine protease is freed by light activation.

A non-limiting list of reactive group reagents corresponding to L-B of R_{11} include
 25 maleimides, N-hydroxysuccinimidyl compounds, imidoesters, 2-iminothiolane, hydrazides, maleic anhydride, and the like, which are available from Pierce Chemical, for example. The foregoing list is merely made for purposes of illustration. Once the reactive group is attached, the L-Z-CINN- X_1A_1 can be attached to a desired molecule (B) such as a targeting mAb using standard conjugation techniques. A non-limiting list of suitable
 30 molecules include antibodies, fragments thereof, single chain binding antibodies and the like, including moieties that are capable of binding/ immobilizing on a particle, for example, Herceptin® (trastuzumab), other monoclonal antibodies, such as those directed to cell surface antigens, murine monoclonal antibodies, proteins, nucleic acids, lectins, lipids, carbohydrates, PAOs, glycosaminoglycans, poly-aspartic acid, poly-L-lysine,

polyvinylpyrrolidone, collagen, peptides, hormones, ligands for receptors, and the like.

Further specific examples are set forth below:

Targeting Agents (including but not limited to the following):

•Any Antibody that targets a cell or tumor cell in some capacity;

- 5 •Monoclonal antibodies (mAbs) with origins from mammals including mice, rats, humans, monkeys, chimeric constructions, etc.;

•Single chain antibodies;

These antibodies can be expressed in bacteria, plants, yeast, animals, mammal milk (mouse, goat, sheep, pig, cow, etc), and animal cell cultures including murine, rat, human,

- 10 hamster, etc.;

•Growth factors both natural and recombinant and peptide fractions of growth factors that bind to receptors on the cell surface (EGF, VEGF, FGF, ILGF-I, ILGF-II, TGF)

•Interferons both natural or recombinant and peptide fractions of interferons that bind to receptors on the cell surface (IFN- α , β , and γ) and interferon agonists;

- 15 •Cytokines, either natural or recombinant, and peptide fractions of cytokines that bind to receptor cell surfaces (IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL10, IL-12, IL-15, TNF, etc);

•Any peptide, natural, recombinant, or synthetic that binds to a cell surface receptor;

•Any hormone either natural or synthetic that binds to a cell surface receptor (estrogen, pro-estrogen);

- 20 •Any metabolite either natural or synthetic that binds to a cell surface receptor (amino acids, sugars, vitamins, nucleotide, nucleoside);

•Any lectin either natural or recombinant that binds to the tumor cell surface;

•Sugar peptides either phosphorylated or non-phosphorylated (MDP, MTP, DDP, DTP)

- 25 or sugar-peptide-lipid targeting agent (monophosphoryl Lipid A, diphosphoryl-Lipid A, DPG-DDP, DPG-DTP, etc.);

•Polyethylene glycol polymers and derivatives (2,000 -> 200,000 Daltons);

•Poly-aspartic or glutamic acids or poly-lysine amino acid polymers or mixtures of these or other amino acids;

- 30 •Inhibitors (peptides or chemical; covalent binding or non-covalent binding) of cell surface enzymes (matrix metalloproteinases inhibitors, tyrosine kinase inhibitors, serine protease inhibitors, casein kinase inhibitors, plasminogen activator inhibitors); and
•glycoseaminoglycans and dextrans.

As will be apparent to those skilled in the art, those targeting agents not specifically mentioned, but falling within the above categories, are also within the scope of the present invention.

Thus, in one aspect of the invention, preferred R_{11} substituents are proteins such as antibodies to biological materials, which can be used to assist in the localizing of potential enzyme activity prior to release by light or other energy source.

D. Carriers and Supports to Immobilize The Z-CINN Derivatives.

Compounds of Formula (I) including Z-CINN core molecule conjugates of inhibited enzymes can be immobilized on a support made of any of a variety of organic or inorganic materials, including synthetic polymers, natural polymers and biosynthetic polymers. These can be in the form of columns, films, membranes, beads, particles, microparticles, and filters.

Immobilization is generally achieved by formation of a covalent, ionic, or affinity bond between a side chain of the core inhibitor molecule (for example 16, N-CINN-AP) and the support material. For example, a support material containing an aldehyde group or reactive acid group could be used. In one example, agarose beads containing free aldehyde groups (Aldehyde-Agarose; Sigma) are added to 16 at pH ~7.0 to form Bead-N-CINN-AP. Excess reagents are washed away and the immobilized inhibitor can then be reacted with a serine protease at pH 6.5 for 12-16 hours at ambient temperature to obtain an immobilized acyl-enzyme, which can be reactivated to give serine protease with light. In another example, agarose beads containing N-hydroxysuccinimidyl groups (Sigma) are added to 16 and processed as above. In another example, utilizing affinity binding, Protein A-Agarose (Sigma or Oncogene) is added to 22 at pH 7, mixed by gentle rocking for 90 min, and separated by centrifugation. The beads are washed several times to remove unbound material and then can be used to bind enzyme as above, and release enzyme activity after exposure to light. These types of binding are within the knowledge of one skilled in the art and can be readily achieved using standard techniques available in the art.

30

III. Cleavage of Acyl Bond

The covalent acyl bond between the cinnamate core molecule and the therapeutic moiety designated in Formula (I) as " X_1A_2 " can be cleaved by hydrolysis, naturally occurring in solution or, at a controllable rate, following energy input. A variety of energy

forms, including light (ultraviolet, visible, or infrared), ultrasound, microwave radiation, radio energy and radioactive decay can be used. Light is most preferred. Light is readily and selectively controllable by variation of the wavelength, duration or energy of the radiation. The preferred energy source is light having a wavelength in the range from 300 to 420 nm, most preferably between 350 nm and 400 nm.

The acyl bond between the enzyme and the enzyme inhibitor can be susceptible to hydrolysis at neutral pH in the dark. The rate of hydrolysis can be modulated by variation of pH. Its half-life can vary from seconds up to several days. The substituted amino group on the cinnamate ring confers stability to the Z-CINN-acyl-enzyme bond that lasts preferably >24 hr at pH 7.0. This is in contrast to the acyl-enzyme bond reported by King (US Patent No. 5,770,699) whose hydrolysis rate is designed to be 20-90 minutes. This stability makes the cinnamate based inhibitors useful for therapeutic applications where the inhibited enzyme needs to be kept inhibited in an aqueous system until activation with light. The Z-CINN-acyl-enzyme can therefore also serve as a prodrug carrier if desired which releases the enzyme in vivo without light activation.

Reactivation of the Z-CINN-X₁-A₂ enzyme is accelerated by light energy. Energy input induces a conformational change in the Z-CINN portion leading to hydrolysis of the acyl-enzyme bond and release of active enzyme. The preferred energy sources have a wavelength in the range from 300 to 420 nm for reactivation times ranging from seconds to minutes. The extent of reactivation depends on such factors as the nature of the inactivated enzyme, the environment, and the type and intensity of the energy source. For example, inactivated thrombin can be readily reactivated by application of white light in a period of about twenty seconds.

The energy source can be introduced via means such as an external visible light source like a focused lamp or laser. A fiber optic catheter that emits light from its tip can deliver light to the target. In other embodiments, the energy source can be introduced in the form of microwave irradiation, radio wave irradiation, ultrasound, radiation of radioactive materials, ultraviolet irradiation, or infrared irradiation.

IV. Methods of Use/ Treatment

The compounds disclosed herein have many applications. One general application of pharmaceutical importance uses the bifunctional aspect of the cinnamate backbone to target an enzyme to some specific structure in the body. These structures could be on tumor cells or extracellular, such as a blood clot. Another use is to inhibit an enzyme,

which is not needed or desired initially in an application, and reactivate it later in the procedure. The inactivated enzyme can be reactivated by applying light, ultrasound, or other energy source to cleave the acyl bond between the inhibited enzyme and the inhibitor. Another application is to increase the stability or circulatory lifetime of the inhibited enzyme in solution. The amount of the compound administered to a mammal in need of such treatment will of course depend upon the needs of the patient and the enzyme or therapeutic agent administered. It is contemplated that the amount administered will be based on known amounts for the unconjugated enzyme and will be sufficient to successfully treat the mammal.

The compounds of the present invention are thus administered as part of pharmaceutically acceptable formulations, e.g. as part of parenteral, i.e. intravenous or oral dosage forms as such are known to those of ordinary skill in the art.


The compounds can be used for purification, particularly those that are immobilized and bind selectively with a target molecule under a given set of conditions, by allowing solutions of the targeted enzyme to equilibrate with inhibitor, then washing away solution before releasing free enzyme following energy input. For example, a compound containing 4-aminoiminophenol at the "A" position could be immobilized and used to isolate thrombin from solution.

The compounds can be used in diagnostic assays. The diagnostic assay may be manual or automated, useful either in laboratories or in the form of a kit.

V. Methods of Synthesis

In another aspect of the invention there are provided methods of preparing the conjugates described herein. Some preferred methods include reacting a compound of

Formula (IV):



(IV)

wherein all variables are as defined with respect to Formula (I) and R'₇ is selected from among H, CH₃ and C₂-C₁₀ alkyls;

with a compound of the Formula (V)




wherein L_1 is a moiety containing a functional group capable of reacting with the NH_2 of Formula (IV);

and B_1 is polymer, a biologically active material such as an mAb or fragment, etc.

5

polymeric support or the like;

under conditions sufficient to form a conjugate corresponding to Formula (VI)



(VI)

wherein L_1' represents the residue of L_1 after the reaction of (IV) with (V).

For purposes of the present invention, "conditions sufficient" shall be understood to

10

include not only those conditions set forth in the Examples below but also those conditions, e.g. temperatures, suitable reagents, solvents, etc. associated with achieving the desired final product.

Some particularly preferred compounds of the present invention are set forth in the following examples and figures. See, for example, compounds 16, 18, 20 and 21.

15

The compounds and methods of use thereof disclosed herein can be further understood by the following non-limiting examples.

VI. Examples

20 Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described below are offered by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled.

25 **Example 1. Synthesis of 3-[2-hydroxy-4-(ethyl)(2-butyloxycarbamidoethyl)-amino]phenyl-2-methyl-2-propenoic acid (13)**

Step 1. Synthesis of (2-ethylaminoethylcarbamic acid, 1,1-dimethylethyl ester) (3)

Refer to Figure 1. A 25g (0.28 mol) quantity of N-ethylethylenediamine (1) was placed in a dried 500 ml rb flask. THF, 250 ml, previously dried by distillation from CaH₂, was added via cannula. The flask was kept under argon and cooled in an ice bath for 20 minutes. A dried addition funnel was charged with 50 ml THF to which was added 29.3 ml (0.13 mol) of di-tert-butyl dicarbonate (2). This mixture was added slowly dropwise to the stirred amine. After addition was complete, the reaction was removed from the ice bath and allowed to stir and reach ambient temperature overnight. The volatiles were removed by rotary evaporation. Saturated NaCl (50 ml) was added, and the result extracted with 4 x 100 ml ethyl acetate. The combined organic fractions were dried over Na₂SO₄ overnight. The drying agent was removed by filtration and the volatiles removed by rotary evaporation to yield 2-ethylaminoethylcarbamic acid, 1,1-dimethylethyl ester (3; 27.55g, 0.15 mol). The material was used without further purification.

Step 2. Synthesis of (2-(ethyl)(3-cyclohex-2-ene-1-one)aminoethylcarbamic acid, 1,1-dimethylethyl ester) (5)

A 200 ml quantity of benzene was added to 27.5g (0.15 mol) 2-ethylaminoethylcarbamic acid, 1,1-dimethylethyl ester (3) in a 500 ml rb flask. The flask was equipped with a Dean-Stark trap and condenser. The mixture was brought to reflux under argon and 18g (1.1 eq.) of 1,3-cyclohexanedione (4) was added all at once. The benzene/water layer was removed from the trap, about 75 ml overall. The trap was rinsed with acetone, dried briefly and quickly replaced and filled with freshly activated 4A molecular sieves and the reaction allowed to proceed overnight. The reaction was cooled to ambient temperature under argon. The volatiles were removed by rotary evaporation to yield a red-orange oil, which may solidify on standing. The material (2-(ethyl)(3-cyclohex-2-ene-1-one)aminoethylcarbamic acid, 1,1-dimethylethyl ester; 5) was used without further purification.

Step 3. Synthesis of (2-(ethyl)(2-hydroxyphenyl)aminoethylcarbamic acid, 1,1-dimethylethyl ester) (6)

To 0.15 mol crude 2-(ethyl)(3-cyclohex-2-ene-1-one)aminoethylcarbamic acid, 1,1-dimethylethyl ester (5) was added 300 ml CH₃CN, freshly distilled from CaH₂. The mixture was brought to reflux under argon and 1.1 eq. mercuric acetate, Hg(AcO)₂, was added in one portion. The suspension was stirred at reflux overnight under argon. Reaction was accompanied by a color change to a claret solution, a violet precipitate and elemental Hg. The reaction was allowed to cool to ambient temperature under argon and was then transferred to centrifuge tubes and centrifuged at 3000 rpm for 10 minutes. The

- supernatant solutions were removed and combined. Additional CH_3CN was added to the tubes; the solid precipitate was resuspended and centrifuged as above. The supernatants were pooled with those previously obtained. The volatiles were removed by rotary evaporation. The dark purple isolate was transferred to a separatory funnel with a minimal volume of chloroform and the resulting solution was washed with 5% NaHCO_3 until the pH of the aqueous layer was about 7. The organic layer was removed and two additional extractions were performed with chloroform, followed by three extractions with ethyl acetate. The organic fractions were combined, dried over Na_2SO_4 and freed of volatiles by rotary evaporation. The crude material obtained in this way can be stored under argon at -80°C for up to several days. A 680g quantity of silica was mixed with 134g decolorizing charcoal, slurried with hexane and poured into a 100 mm glass column with about 190 mm of packing. The crude isolate was loaded in a minimum volume of chloroform (3 x 50ml). The elution was performed at 7-10 psi with ethyl acetate/hexane: 1.5 L, 20%; 1.5L, 30%; 1.5L, 40%; 1L, 50%. Fractions were pooled and volatiles removed by rotary evaporation to leave a slightly yellowish oil. The material (2-(ethyl)(2-hydroxyphenyl)aminoethylcarbamic acid; 1,1-dimethylethyl ester; 6) exhibits behavior consistent with sensitivity to air and light at this point and must be handled accordingly. It was successfully stored at -20°C under argon for a period of one month. Yield: 8.78g (21% for two steps).
- Step 4. Synthesis of (3-(ethyl)(2-acetylaminoethyl)aminophenol)(8)**
- Degassed ethyl acetate (75 ml) was added to 5.5g (19.6 mmol) 2-(ethyl)(2-hydroxyphenyl)aminoethylcarbamic acid, 1,1-dimethylethyl ester (6). To this was added 50 ml of 3N HCl and the reaction was stirred at ambient temperature overnight. TLC analysis was done using 30% EtOAc/hexanes. The reaction mixture was transferred to a separatory funnel and extracted with 2 x 40 ml 1N HCl. The pooled aqueous layers were placed on the rotary evaporator and the volatiles removed. Water was added (about 50 ml) and removed by rotary evaporation, 2 cycles. The resulting oil was frozen in dry ice and placed under vacuum, then allowed to warm naturally to remove all volatiles. The resulting crispy material (7) was used without further purification, and kept free of air and moisture. Dry pyridine (75 ml) was added to the isolate by cannula. The flask was swirled to dissolve the material. Acetic anhydride (1.2 eq.) was added via syringe and the reaction was stirred under argon at ambient temperature. TLC analysis was performed with 10% methanol/chloroform. At the completion of the reaction, the pyridine was removed by rotary evaporation and the residue treated with 100 ml of 10% NaOH by

stirring overnight under argon. The reaction was adjusted to pH 9 with 1N HCl and extracted with 4 x 75 ml ethyl acetate. The combined organic layers were washed with water, then brine (once each), pre-dried with Na₂SO₄, then dried overnight with MgSO₄. The solution was collected by filtration and the volatiles were removed by rotary

5 evaporation. The material was purified by flash chromatography on silica: loaded with 5% methanol/chloroform; eluted with 150 ml each of 3.3, 6.6, and 9.9% solvent as above to give 3-(ethyl)(2-acetylaminoethyl)aminophenol (8). Yield was 4.2g (96%).

Step 5. Synthesis of (2-hydroxy-4-(ethyl)(2-acetylaminoethyl)aminobenzaldehyde) (9)

A scrupulously dry flask was charged with anhydrous DMF (15 ml) and cooled in
10 ice under argon. Phosphorus oxychloride, POCl₃, 1.5 ml (1.15 eq) was added slowly dropwise to the stirred solution. After the addition was complete, the flask was removed from the ice bath and allowed to reach ambient temperature. This was accompanied by a color change to yellow. The flask was heated in a 40°C bath for one hour. It was removed from the heat, and once at ambient temperature, a DMF solution of 3-(ethyl)(2-
15 acetylaminoethyl)aminophenol (8; 3.2g, 14 mmol) was added via addition funnel. After the addition was complete, the flask was returned to the 40°C bath and stirred overnight. The reaction was cooled to 0°C and 25 ml 10% NaOH was added, the mixture was allowed to warm to ambient temperature and was then placed in a 40° C bath. Additional NaOH was added to achieve pH >10. Water (50 ml) was added. After 15 minutes the flask was
20 removed from the bath and the pH adjusted to 5-5.5 with 1N HCl. The mixture was transferred to a separatory funnel and extracted with 4 x 100 ml ethyl acetate. The pooled organic fractions were washed with water (1 x 75 ml), predried with Na₂SO₄, and then dried with MgSO₄ overnight. The solution was collected by filtration and the volatiles removed by rotary evaporation to yield the crude product (2.9 g). The material was
25 purified by flash chromatography on silica, 45 x 150 mm column, eluting with 200 ml each of 2.5%, 5%, and 10% methanol/chloroform. Purified yield of (2-hydroxy-4-(ethyl)(2-acetylaminoethyl)aminobenzaldehyde) (9) was 1.52 g (45%).

Step 6. Synthesis of (3-[2-hydroxy-4-(ethyl)(2-acetylaminoethyl)amino]phenyl-2-methylpropenoic acid, ethyl ester) (11)

30 2-hydroxy-4-(ethyl)(2-acetylaminoethyl)aminobenzaldehyde (9; 1.58g, 6.3 mmol) was placed in a flask with degassed benzene (50 ml). Carbethoxy(ethylidene)triphenylphosphorane (10; 1.05 eq.) was added and the reaction stirred overnight under argon. The reaction was freed of volatiles by rotary evaporation, then purified by flash chromatography (silica, 95mm x 45mm), eluting with 250 ml 2.5%;

150 ml 5%; and 150 ml 10%; all methanol/chloroform to give 3-[2-hydroxy-4-(ethyl)(2-acetylaminoethyl)amino]phenyl-2-methylpropenoic acid, ethyl ester (11). The yield was quantitative.

Step 7. Synthesis of (3-[2-hydroxy-4-(ethyl)(2-tert-butyloxycarbamidoethyl)-amino]phenyl-2-methylpropenoic acid, ethyl ester) (12)

5 3-[2-hydroxy-4-(ethyl)(2-acetylaminoethyl)amino]phenyl-2-methylpropenoic acid, ethyl ester (11; 2.0g, 6 mmol) was dissolved in anhydrous THF (30 ml). Di-*t*-butyldicarbonate (2; 3 eq) and dimethylaminopyridine (0.1 eq) were added and the reaction was heated to 75°C. TLC analysis (silica, 5% methanol/chloroform) was used to
10 gauge reaction completion; after 3.5 hours the reaction was determined to be finished. The solution was allowed to cool to ambient temperature, the volume of the reaction was doubled with anhydrous methanol, and hydrazine (3 eq.) was added dropwise via syringe. The reaction was allowed to stir under argon overnight. The solution was poured into 100 ml CH₂Cl₂ in a separatory funnel, and water (100 ml) was added. The mixture was shaken
15 and allowed to separate; the water layer was adjusted to pH 4.5 by judicious addition of 1N HCl, shaking well after each addition. The aqueous layer was cloudy but colorless. After thorough washing, the layers were separated and the organic layer washed with aqueous NaHCO₃ at pH 7.8-8.0. The organic layer was dried by stirring overnight over MgSO₄. The solution was collected by filtration and the volatiles were removed by rotary
20 evaporation. The material was purified by flash chromatography on silica, using 400 ml 15% and 300 ml 30% ethyl acetate/hexane to give 3-[2-hydroxy-4-(ethyl)(2-tert-butyloxycarbamidoethyl)amino]phenyl-2-methylpropenoic acid, ethyl ester (12). The yield was 1.7g (72%).

Step 8. Synthesis of 3-[2-hydroxy-4-(ethyl)(2-tert-butyloxycarbamidoethyl)-amino]phenyl-2-methylpropenoic acid (13)

25 3-[2-hydroxy-4-(ethyl)(2-tert-butyloxycarbamidoethyl)amino]phenyl-2-methylpropenoic acid, ethyl ester (12; 759mg, 1.9 mmol) was taken up in ethanol (10 ml). A 5 ml quantity of 10% w/v LiOH was added and the reaction was heated to 40°C under argon. The reaction was complete after 6 hours, determined by TLC. The mixture was
30 allowed to cool to ambient temperature, the pH adjusted to 5-6 with 1N HCl, and extracted into ether (4 x 50 ml); the combined organic layers were washed with saturated NH₄⁺Cl⁻ and dried over CaSO₄ overnight. The solution was collected by filtration and the volatiles were removed by rotary evaporation to give 3-[2-hydroxy-4-(ethyl)(2-tert-

butyloxycarbamidoethyl)amino]phenyl-2-methylpropenoic acid (13). The material was used without further purification. Yield was 509mg (74%).

Example 2. Preparation of 3-[2-hydroxy-4-(ethyl)aminoethyl)amino]phenyl-2-methylpropenoic acid, 4-aminoiminophenyl ester, HCl salt (16; N-CINN-AP)

5 **Step 1. Synthesis of 3-[2-hydroxy-4-(ethyl)(2-tert-butyloxycarbamidoethyl)amino]-phenyl-2-methylpropenoic acid, 4-aminoiminophenyl ester, HCl salt (15; tBOC-N-CINN-AP)**

Refer to Figure 2. 3-[2-hydroxy-4-(ethyl)(2-tert-butyloxycarbamidoethyl)-amino]phenyl-2-methylpropenoic acid (13) (509 mg; 1.4 mmol) was dissolved in
10 anhydrous pyridine. Dicyclohexylcarbodiimide (1.2 eq.), 4-aminoiminomethylphenol hydrochloride (14; 1.1 eq.) and dimethylaminopyridine (0.05 eq.) were added and stirred at ambient temperature under argon for 12-16 hr. The reaction mixture was passed through a medium porosity sintered glass filter, and freed of volatiles by rotary evaporation. The isolate was taken up in 5% methanol/chloroform, filtered, and freed of
15 volatiles as above. The crude material, 15, was purified by flash chromatography on silica, loading with 3 ml 5% and eluting with 100 ml 5%; 75 ml 10%; 75 ml 15%; and 200 ml 20%; all methanol/chloroform. Yield was 448mg (66%).

Step 2. Synthesis of 3-[2-hydroxy-4-(ethyl)aminoethyl)amino]phenyl-2-methylpropenoic acid, 4-aminoiminophenyl ester, HCl salt (16; N-CINN-AP)

20 3-[2-hydroxy-4-(ethyl)(2-tert-butyloxycarbamidoethyl)amino]phenyl-2-methylpropenoic acid, 4-aminoiminophenyl ester, HCl salt (15; 12 mg, 25 μ mol) was suspended in 1.2 ml of 3N HCl and mixed by inversion for 3 hours at ambient temperature to release the t-BOC group and generate 3-[2-hydroxy-4-(ethyl)aminoethyl)amino]phenyl-2-methylpropenoic acid, 4-aminoiminophenyl ester, HCl salt
25 (16; N-CINN-AP). After 3 hours, the reaction was neutralized with 3N NaOH to give a final pH of 7.0. This material was used immediately and was not characterized further except that it was now soluble in aqueous solution.

Example 3. Preparation of mPEG₅₀₀₀-N-CINN-AP (18)

30 Refer to Figures 3 and 4. mPEG₅₀₀₀-SS (2M4R0H01; Shearwater Polymers) was used as follows. 3-[2-hydroxy-4-(ethyl)aminoethyl)amino]phenyl-2-methylpropenoic acid, 4-aminoiminophenyl ester, HCl salt (16; 2.5 mg, 6.5 μ mol; pH 7.0-7.4) in 1 ml of 10 mM sodium phosphate buffer, pH 7.2, was added to dry mPEG₅₀₀₀-SS (17; 38.3 mg, 7.5 μ mol). After 1 hour at ambient temperature, the reaction was stopped by the addition of

75 μ M glycylglycine in 100 μ l of water. This material (mPEG₅₀₀₀-NH-CINN-AP; 18) was used within 3 hours for inhibition reactions with serine proteases such as thrombin and tPA.

5 Example 4. Inhibition of Thrombin with R-NH-CINN-AP Derivatives.

R is -tBOC, -H₂⁺, or mPEG₅₀₀₀. Human thrombin was modified with R-NH-CINN-AP inhibitors as follows: Thrombin (600 units; ~ 3.5 nmol) in 50 mM sodium citrate buffer, 3 mM CaCl₂, 7% kollidon, pH 7.0, was reacted with 3.5-35 nmol of the R-N-CINN-AP inhibitor for 12-18 hours at ambient temperature in the dark. Thrombin activities were measured under red light ("dark") and after exposure to 10 seconds of blue light from a dental light and were assayed using S-2238 (diaPharma) at 30°C. Percentage inhibition was calculated as 100 - dark activity. Dark activity was calculated as [activity_{dark} ÷ activity_{control}] X 100. Light reactivation was calculated as activity_{light} ÷ activity_{control}. The following are typical results.

15

Table 1. Inhibition and Reactivation of R-N-CINN-Acyl-Thrombin^a

Inhibitor	Inhibitor Excess	Inhibition after 16 Hrs Incubation (%)	Light Reactivation (%)
None	0	0	NP
CINN-AP	1X	53	NP
CINN-AP	5X	99	NP
CINN-AP	10X	99	60
#15	1X	94	NP
#15	5X	98	NP
#15	10X	97	72
#16	1X	21	NP
#16	5X	70	NP
#16	10X	96	64
#18	2X	51	NP
#18	10X	98	69

^aThrombin activity was assayed as described in Chromogenic Substrates (Chromogenix AB Information Manual) using S-2238 (H-D-Phe-Pip-arg-pNA•2HCl) as the synthetic substrate.

20 #15 = tBOC-N-CINN-AP
 #16 = NH₃-CINN-AP
 #18 = mPEG₅₀₀₀-N-CINN-AP
 NP = Assay not performed

25 The R-NH-CINN-AP compounds inhibit thrombin, as does the parent compound CINN-AP. All of the acyl-thrombins can be reactivated with blue light, indicating that the

modifications to create R-NH-CINN-AP do not affect the ability of the molecule to isomerize and release free enzyme from the acyl-enzyme. The NH₃-CINN-AP, which contains a positive charge, appears to be a less effective inhibitor at lower molar ratios. The mPEG₅₀₀₀-NH-CINN-AP, which contains a 5000 dalton linear mPEG molecule, requires higher concentrations to effect thrombin inhibition. These results show that one can alter other portions of the molecule without losing the effects of a light reversible, covalent inhibitor. Light reactivation remains essentially the same, indicating that the R-NH-CINN-acyl-thrombin molecule absorbs blue light energy and can isomerize to give the *cis* configuration and release active thrombin.

Example 5. Inhibition of Tissue Plasminogen activator (tPA) with R_x-NH-CINN-AP Derivatives.

R_x is -tBOC, -H₂⁺, or mPEG₅₀₀₀. The procedures of Example 4 was repeated except that Human tPA (Activase, Genentech Inc. Lot # E9034A) was modified with R_x-NH-CINN-AP inhibitors in place of thrombin. The reactions were carried out as follows: tPA (0.83 nmol) in 10 mM tris-HCl, 130 mM NaCl, pH 7.4 was reacted with 0.83-8.3 nmoles of the R-NH-CINN-AP inhibitors for 12-18 hours at ambient temperature in the dark. Activities were calculated as for Example 4, using S-2288 (diaPharma). The results are provided below.

Table 2. Inhibition and reactivation of R_x-N-CINN-tPA^a

Inhibitor	Inhibitor Excess	Inhibition after 16 hrs incubation (%)	Light Reactivation (%)
None	0	0	NP
CINN-AP	5X	95	NP
CINN-AP	10X	95	82
#15	5X	95	NP
#15	10X	95	87
#16	5X	78	NP
#16	10X	88	78

^a tPA activity was assayed as described in Chromogenic Substrates (Chromogenix AB Information Manual) using S-2288 (H-D-Ile-Pro-Arg-pNA•2HCl) as the synthetic substrate

#15 = tBOC-N-CINN-AP

#16 = NH₃-CINN-AP

NP = Assay not performed

The R_x-NH-CINN-AP compounds inhibit tPA, as does the parent compound CINN-AP. All of the acyl-tPAs can be reactivated with blue light, indicating that the modifications to create R-NH-CINN-AP do not affect the ability of the molecule to

isomerize and release free enzyme from the acyl-enzyme. The NH₃-CINN-AP, which contains a positive charge, appears to be a less effective inhibitor at lower molar ratios. These results show that other portions of the molecule can be altered without losing the effects of a light reversible, covalent inhibitor. Light reactivation remains essentially the same, indicating that the R-NH-CINN-acyl-tPA molecule absorbs blue light energy and can isomerize to give the *cis* configuration and release active tPA.

Example 6. Preparation of N-CINN-AP with Maleimide Linker (20)

N-CINN-AP (16; 10 μmoles, 1 ml) was reacted with 11.4 μmoles of SMPH (succinimidyl-[β-maleimidopropionamido] hexanoate; Pierce #22363) in 0.5 ml DMSO and 0.5 ml 200 mM sodium phosphate, pH 7.4 for 2 hours at ambient temperature. After 2 hours, the reaction was placed on ice, and unreacted NHS was quenched with 0.05 ml of 100 mM glycylglycine. The final product (20) was used within 2 hours, and was not further characterized.

Example 7. Preparation of Antibody-Linker-N-CINN-AP (21)

A monoclonal antibody, Herceptin® (trastuzumab, Genentech) was prepared at 5 mg/ml in phosphate-NaCl buffer, pH 7. SATA (N-succinimidyl S-acetylthioacetate; Pierce) was dissolved in DMSO and a 20-fold molar excess was added to the protein and reacted for 1 hour. Hydroxylamine HCl (Pierce; 0.5 M, neutralized to pH 7.0) was added for 60-90 minutes, and antibody was separated from low molecular weight material by chromatography on D-SALT Polyacrylamide desalting columns (Pierce) equilibrated in phosphate-NaCl buffer. Incorporation of thiols into antibody (antibody-SH) was determined using Ellman's reagent (Pierce). 20, at a 2-fold molar excess to incorporated – SH groups, was added to Antibody-SH for 2 hours at pH 7.4 and ambient temperature to give 21. Antibody-containing fractions (21) were separated from excess 20 on a D-SALT column as above.

Example 8. Preparation of Antibody-Linker-N-CINN-Enzyme (22)

21 was mixed with thrombin (400 U/ml; Enzyme Research Labs) at pH 7.0 for 12-16 hours at ambient temperature to form 22, and 22 was separated from free enzyme by binding to Protein A-Agarose beads (Oncogene) which had been washed 3 times with phosphate-NaCl buffer. After gentle mixing for 90 min, the beads were washed by centrifugation 3 times and supernatants containing unbound enzyme were removed. Beads were suspended in assay buffer at pH 8.3 containing 0.5 mM S-2238 (Chromogenix), and

assayed for 6 min at 30°C. Ten seconds of blue light from a ProLite dental light was then directed to the wells, and assays were continued for another 6 min. The rate of S-2238 cleavage increased 7-8 fold on exposure to light, indicating release of free thrombin into solution.

5

Example 9. Preparation of Antibody-PEG-N-CINN-AP (23)

N-CINN-AP (16, 2.4 μ moles, 1.0 ml) in 50 mM sodium phosphate buffer, pH 7.0, was reacted with 4.8 μ moles of PEG₃₄₀₀-(SPA)₂ (PEG succinimidyl propionate; Shearwater Polymers, Inc) in 1 ml WFI at ambient temperature for 1 hour. The SPA-PEG-N-CINN-AP was immediately reacted with goat anti-rabbit IgG (Calbiochem, 401311) as follows: 10.6 nmoles of SPA-PEG-N-CINN-AP was reacted with 2.6 nmoles antibody for 2 hours at pH 7.0 at ambient temperature. The reaction mixture was placed on ice and excess SPA was quenched with 0.05 ml of 100 mM glycylglycine. The product was used within two hours, and not further characterized.

15

Example 10. Preparation and Assay of Antibody-PEG-N-CINN-Acyl-Thrombin (24)

Antibody-PEG-N-CINN-AP (23) was mixed with thrombin (400 U/ml; Enzyme Research Labs) at pH 7.0 for 16 hours at ambient temperature. The antibody-acyl-enzyme complex was then tested for its ability to bind to its antigen and release active enzyme. Rabbit IgG (Calbiochem) was bound to polystyrene plates following standard procedures. The plates were then blocked with PBS/Tween 20/BSA (Sigma) for 2 hours, and washed extensively with PBS/Tween 20 to remove excess protein. Goat anti-rabbit antibody-PEG-N-CINN-Acyl-Thrombin (24) was then allowed to bind to the rabbit IgG on the plate for 2 hours at ambient temperature. The plates were then washed 5 times with PBS/Tween 20 to remove any unbound antibody-thrombin or free thrombin. The plates were assayed for thrombin activity using the S-2238 chromogenic assay described in Example 8. The rate of S-2238 cleavage increased 5 fold after 10 seconds of light activation indicating release of free thrombin into solution.

20

25

30

Examples 11-13 Murine mAbs As Targeting Agents

The following examples provide details regarding the preparation of specific murine mAb-AZ-CINN conjugates and their utility as targeting agents for therapeutic indications.

35

Example 11. Lung Cancer**A. Preparation of Antibody-Linker-CINN-Enzyme and Its Use**

5 N-CINN-AP (16, 2.4 μ moles, 1.0 ml) in 50 mM sodium phosphate buffer pH 7.0 is reacted with 4.8 μ moles of NHS-PEG₃₄₀₀-MAL (Shearwater Polymers, Inc. NHS-Mal-3400) dissolved in DMSO. The reaction mixture is rocked for two hours at ambient temperature. The reaction is stopped by the addition of 100 μ l of 100 mM glycylglycine. The Mal-PEG₃₄₀₀-N-CINN-AP is used within two hours and not further characterized.

B. Preparation of Antibody-PEG-N-CINN-Acyl-Thrombin

10 A murine monoclonal antibody, BR110 (Hellstrom et.al. US Patent 5,840,854) is diluted to 5 mg/ml in phosphate-NaCl buffer, pH 7. SATA (N-succinimidyl S-acetylthioacetate; Pierce) is dissolved in DMSO and a 20-fold molar excess is added to the protein and reacted for 1 hour. Hydroxylamine HCl (Sigma; 0.5 M, neutralized to pH 7.0) is added for 60-90 minutes, and antibody is separated from low molecular weight material by
15 chromatography on D-SALT Polyacrylamide desalting columns (Pierce) equilibrated in phosphate-NaCl buffer. Incorporation of thiols into the murine antibody (antibody-SH) is determined using Ellman's reagent (Pierce). Mal-PEG₃₄₀₀-N-CINN-AP at a 2-fold molar excess to incorporated -SH groups, is added to BR110-mAb-SH for 2 hours at pH 7.4 and ambient temperature to give BR110-mAb-S- Mal-PEG₃₄₀₀-N-CINN-AP. Antibody-
20 containing fractions (BR110-mAb-S- Mal-PEG₃₄₀₀-N-CINN-AP) are separated from excess Mal-PEG₃₄₀₀-N-CINN-AP on a D-SALT column (Pierce) buffered with 50 mM sodium phosphate pH 6.5 and 150 mM NaCl. The antibody-drug (BR110-mAb-S- Mal-PEG₃₄₀₀-N-CINN-AP) is concentrated to 5 mg/ml protein using Centricon YM-30 membranes and sterile filtered (0.22 μ m filter; Gelman Sciences, Supor Acrodisc 25).
25 BR110-mAb-S-Mal -PEG-N-CINN-AP is mixed with thrombin (400 U/ml; Enzyme Research Labs) at pH 7.0 for 16 hours at ambient temperature. The antibody-acyl-enzyme complex is then separated from free thrombin and concentrated to 5 mg/ml protein using Centricon YM-100 membranes and sterile filtered (0.22 μ m filter; Gelman Sciences, Supor Acrodisc 25).

30 **C. Treatment of Human Lung Carcinoma with BR110-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin**

MAB BR100 is a murine mAb to a 66-kDa glycoprotein that is found on the cell surface of human lung, colon and breast tumor cells. Human lung carcinoma cell line H2987 is grown in cell culture as described (U.S. Patent No.5,840,854). 1×10^7 cells/

mouse are implanted s.c. into a nude mice. When the tumors reach 150-200 mm³ in size, BR110-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin is used to treat the tumor as follows; 1.0 mg of BR110-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin (1 mg mAb protein with ~ 27 nmoles thrombin covalently attached) is injected i.v. via tail vein into mice with the tumor. After 6 hours, mice that are given the BR110-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin are anesthetized with a Ketamine/Rompun mixture. A 16G needle is used to create a hole in the skin and a 1 mm fiber optic probe is inserted up to the tumor. The tumor is exposed to white light for 5 minutes with a power of about 0.5 mW/cm². Other treatment groups do not receive light treatment. All mice are kept under "red lights" to prevent light activation of the drug, for 72 hours post injection, then returned to normal light/dark cycle. Tumors in groups that receive light treatment shrink immediately (>50% in 7 days) and are necrotic. Tumors that are treated with BR110-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin but do not receive light stimulation shrank at a much slower pace and are smaller than 100mm³ at 28 days. Groups that receive antibody and free drug (unlinked) show tumor growth, and by 28 days the mice either die or have tumors > 800 mm³.

Example 12. Colon Cancer:

- A. Preparation of Antibody-Linker-CINN-Acyl-Thrombin
 Mal-PEG₃₄₀₀-N-CINN-AP is prepared as described above in Example 11A.
- B. Preparation of Antibody-PEG-N-CINN-Acyl-Thrombin
 The procedure of Example 11B is repeated except that MAb A7, a murine mAb to a 45-kDa glycoprotein found on human colon cancer cells, is used in place of the BR110-mAb to produce A7-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin.
- C. Treatment of Human Colon Carcinoma with A7-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin
 Human colon carcinoma cell line LS-180 is grown in cell culture as described (Kinuya et. al. 2001. J. Nucl. Med. 42:596-600). 1 x 10⁷ cells/mouse are implanted s.c. into nude mice. When the tumors reach 150-200 mm³ in size, A7-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin is used to treat the tumor as follows; 1.0 mg of A7-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin (1 mg mAb protein with ~ 27 nmoles of thrombin covalently attached) is injected i.v. via tail vein into mice with the tumor. After 6 hours, mice that are given the A7-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin are anesthetized with a Ketamine/Rompun mixture. A 16G needle is used to create a hole in

the skin and a 1 mm fiber optic probe is inserted up to the tumor. The tumor is exposed to white light for 5 minutes with a power of about 0.5 mW/cm². Other treatment groups do not receive light treatment. All mice are kept under "red lights" to prevent light activation of the drug, for 72 hours post injection, then returned to normal light/dark cycle. Tumors in groups that receive light treatment shrink immediately (>50% in 7 days) and are necrotic. Tumors that are treated with A7-mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin but do not receive light stimulation shrink at a much slower pace and are smaller than 100mm³ at 28 days. Groups that receive antibody and free drug (unlinked) show tumor growth, and by 28 days the mice either die or have tumors > 800 mm³.

10

Example 13. Breast Cancer

A. Preparation of Antibody-Linker-CINN-Acyl-Thrombin

Mal-PEG₃₄₀₀-N-CINN-AP is prepared as described above.

B. Preparation of Antibody-PEG-N-CINN-Acyl-Thrombin

The procedure of Example 11B is repeated except that MAb NR-LU-10, a murine mAb to a pancarcinoma glycoprotein found on human breast cancer cells, is used in place of the BR110-mAb to produce NR-LU-10 S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin.

C. Treatment of Human Breast Carcinoma with NR-LU-10 mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin

Human breast cancer xenografts are prepared as described (Burak et.al. 1998. Nucl. Med. Biol. 25:633-637). Xenografts are implanted s.c. into nude mice. When the tumor reaches 150-200 mm³ in size, NR-LU-10 mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin is used to treat the tumor as follows; 1.0 mg of NR-LU-10 mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin (1 mg mAb protein with ~ 27 nmoles of thrombin covalently attached) is injected i.v. via tail vein into mice with the tumor. After 6 hours, mice that are given the NR-LU-10 mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin are anesthetized with a Ketamine/Rompun mixture. A 16G needle is used to create a hole in the skin and a 1 mm fiber optic probe is inserted up to the tumor. The tumor is exposed to white light for 5 minutes with a power of about 0.5 mW/cm². Other treatment groups do not receive light treatment. All mice are kept under "red lights" to prevent light activation of the drug, for 72 hours post injection, then returned to normal light/dark cycle. Tumors in groups that receive light treatment shrink immediately (>50% in 7 days) and are necrotic. Tumors that are treated with NR-LU-10 mAb-S- Mal-PEG₃₄₀₀-N-CINN-Acyl-Thrombin but do not receive light stimulation shrink at a much slower pace and are


smaller than 100mm^3 at 28 days. Groups that receive antibody and free drug (unlinked) show tumor growth, and by 28 days the mice either die or have tumors $> 800\text{mm}^3$.

The various publications, patents, patent applications and published applications mentioned in this application are hereby incorporated by reference herein.

5

WE CLAIM:

1. A compound comprising the formula:




wherein:

- 5 **R₁ and R₂ are individually selected from the group consisting of H, CH₃, C₂-C₁₀ alkyls, C₂-C₁₀ alkenyls or C₂-C₁₀ alkynyls, each of which can be substituted or unsubstituted; straight or branched, C₂-C₁₀ heteroalkyls, C₂-C₁₀ heteroalkenyls or C₂-C₁₀ heteroalkynyls and -(CR₁₅R₁₆)_p-D;**

- 10 **wherein: R₁₅ and R₁₆ are individually selected from the group consisting of H, CH₃, C₂-C₁₀ alkyls, C₂-C₁₀ alkenyls or C₂-C₁₀ alkynyls, each of which can be substituted or unsubstituted; straight or branched; and C₂-C₁₀ heteroalkyls, C₂-C₁₀ heteroalkenyls or C₂-C₁₀ heteroalkynyls; p is a positive integer from 1 to about 12;**

D is selected from among -SH, -OH, X₂, -CN, -OR₁₉, NHR₂₀,



15

wherein:

- R₁₇ is H, CH₃ or X₃;**
R₁₈ is H, a C₁₋₄ alkyl or benzyl;
R₁₉ is H, a C₁₋₄ alkyl, X₂ or benzyl;

20 **R₂₀ is H, a C₁₋₁₀ alkyl or -C(O)R₂₁,**
wherein R₂₁ is H, a C₁₋₄ alkyl or alkoxy, t-butoxy or benzyloxy;
X₂ and X₃ are independently selected halogens;

R₃ is H, CH₃, or -C(=O)(CR₁₅R₁₆)_w-D,


- 25 **where w is 0 or an integer from 1 to about 12, and D is H or as described for R₁ and R₂**

J is O, NH or S;

R₄, R₅, and R₆ are independently selected from the group consisting of H, CH₃, C₂-C₁₀ alkyls, C₂-C₁₀ alkenyls or C₂-C₁₀ alkynyls, each of which can be substituted or unsubstituted; straight or branched; C₂-C₁₀ heteroalkyls, heteroalkenyls or heteroalkynyls and halogens;

5

Z is NR₇R₈ or



wherein R₇ is selected from among H, CH₃, C₂-C₁₀ alkyls, alkenyls or alkynyls which can be substituted or unsubstituted; straight or branched; C₂-C₁₀ heteroalkyls, heteroalkenyls or heteroalkynyls, or -(CR₂₃R₂₄)_q-aryl, or R₈,

10 wherein R₂₃ and R₂₄ are independently selected from the group consisting of H and C₁-C₁₀ alkyls;

q is an integer from 1 to about 6;

R₈ is selected from the group consisting of (CR₉R₁₀)_n-NR₂₂-R₁₁, (CR₉R₁₀)_n-CH₂-NHC(O)R₂₆ and (CR₉R₁₀)_n-CH₂-E;

15 wherein R₉ and R₁₀ are independently selected from the group consisting of H, CH₃, C₂-C₁₀ alkyls, C₂-C₁₀ alkenyls or C₂-C₁₀ alkynyls, each of which can be substituted or unsubstituted; straight or branched; C₂-C₁₀ heteroalkyls, C₂-C₁₀ heteroalkenyls or C₂-C₁₀ heteroalkynyls and halogens;

R₂₆ is H, CH₃, O-t-butyl, O-benzyl;

20 E is OH, SH or O-C(O)R₂₇,

wherein R₂₇ is a C₁-C₆ alkyl, benzyl or phenyl;

R₂₂ is H or CH₃;

n is a positive integer from 1 to about 10;

R₁₁ is H or -L-B,

25 wherein L is a linker; and

B is a first active moiety, reactive group moiety or a polymer;

R₂₅ is H, -C(O)-R₂₈ or -C(O)-O-R₂₉,

wherein R₂₈ is a C₁-C₆ alkyl or benzyl; and R₂₉ is CH₃, t-butyl or benzyl;

X₁ is O, NH, or S; and

30 A is H or a second active moiety.


2. The compound of claim 1, wherein Z is NR₇R₈.

3. The compound of claim 2, wherein R_8 is $-\text{CH}_2-\text{CH}_2-\text{NH}_2$.
4. The compound of claim 2, wherein R_8 is $(\text{CR}_9\text{R}_{10})_n-\text{NR}_{22}-\text{R}_{11}$.
- 5 5. The compound of claim 1, wherein L-B comprises a maleimidyl or an N-hydroxysuccinimidyl group.
6. The compound of claim 4, wherein R_{11} comprises a polyalkylene oxide residue.
- 10 7. The compound of claim 6, wherein said polyalkylene oxide residue is a polyethylene glycol.
8. The compound of claim 7, wherein said polyethylene glycol has a number average molecular weight of from about 2,000 to about 200,000 daltons.
- 15 9. The compound of claim 4, wherein R_{11} comprises a member of the group consisting of collagen, glycosaminoglycan, poly(-aspartic acid), poly(-L-lysine), poly(-lactic acid), poly-N-vinylpyrrolidone and copolymers of poly(-lactic acid) and poly(-glycolic acid).
- 20 10. The compound of claim 1, wherein R_1 , R_2 , R_3 , R_4 , R_5 , and R_6 are independently selected from the group consisting of H, CH_3 and CH_3CH_2 .
- 25 11. The compound of claim 4, wherein R_7 is CH_3CH_2 ; R_8 is $-(\text{CR}_9\text{R}_{10})_n-\text{NR}_{22}-\text{R}_{11}$; and R_9 and R_{10} are H; n is 2; and X_1 is O, S or NH.
12. The compound of claim 4, wherein R_7 is CH_3CH_2 ; R_8 is $-(\text{CR}_9\text{R}_{10})_n-\text{NR}_{22}-\text{R}_{11}$ and R_9 and R_{10} are H.
- 30 13. The compound of claim 1, wherein said second active moiety comprises a member of the group consisting of X_1A_1 or X_1A_2 wherein
 X_1A_1 is a substrate or substrate analog selected from the group consisting of amino acids, amino acid derivatives, peptides, peptide derivatives and substrates or substrate

analogues for serine proteases, cysteine proteases, esterases, lipases, or other enzymes containing an active site serine or cysteine; and

X_1A_2 is an enzyme.

- 5 14. The compound of claim 13, wherein X_1A_1 is a moiety of the formula



wherein R_{12} and R_{13} are independently H or electron donating or electron withdrawing groups and W is CH or N.

- 10 15. The compound of claim 13, wherein A_2 is an enzyme selected from the group consisting of serine proteases, cysteine proteases, esterases, lipases and enzymes containing an active-site serine or cysteine.

16. The compound of claim 14, wherein J is O, R_2 is H, R_7 is CH_3CH_2 ; R_8 is
15 $-(CR_9R_{10})_n-NR_{22}-R_{11}$, R_9 and R_{10} are H, and n is 2.

17. The compound of claim 15, wherein X_1A_2 is an enzyme having an active-site serine or cysteine.



- 20 18. The compound of claim 11, wherein X_1A_2 is a blood coagulation factor.

19. The compound of claim 11, wherein the enzyme is selected from the group consisting of plasmins, urokinases, and tissue plasminogen activators.


- 25 20. The compound of claim 13, wherein X_1A_1 is an amino acid, peptide, or substrate or substrate analog capable of interacting with an enzyme.


21. The compound of claim 20, wherein said amino acid is selected from the group consisting of isoleucine, phenylalanine, tyrosine, lysine, arginine, aspartate, glutamate,
30 glutamine and asparagine.

22. A compound of claim 1 selected from the group consisting of:




5





and



wherein

PEG is a polyethylene glycol having a molecular weight of from about 2,000 to about

5 200,000; and

mAb is a monoclonal antibody.

23. The compound of claim 22, wherein said monoclonal antibody is trastuzumab.

10 24. The compound of claim 1, wherein L-B comprises a maleimidyl or an
N-hydroxysuccinimidyl group.

25. A pharmaceutically acceptable salt of the compound of claim 1.

15

26. A method of treatment, comprising:
administering to a mammal in need of such treatment an effective amount of a compound of claim 1, where B is a first active moiety.

5 27. The method of claim 26, further comprising exposing the compound of claim 1 to an energy source after administration to said mammal.


28. The method of claim 27, wherein the energy source is white light having a wavelength in the range from 340 to 700 nm.

10

29. The method of claim 27, wherein the energy source is white light having a wavelength in the range from 350- 420 nm.

30. The method of claim 27, wherein the energy source is selected from the group
15 consisting of microwave, ultrasound, radio energy, gamma radiation, radioactivity, ultraviolet light and infrared light.

31. A method of preparing a conjugate, comprising:
reacting a compound of Formula (IV)



20 (IV)

wherein:


R₁ and R₂ are individually selected from the group consisting of H, CH₃, C₂-C₁₀ alkyls, C₂-C₁₀ alkenyls or C₂-C₁₀ alkynyls, each of which can be substituted or unsubstituted; straight or branched, C₂-C₁₀ heteroalkyls, C₂-C₁₀ heteroalkenyls or

25 C₂-C₁₀ heteroalkynyls and -(CR₁₅R₁₆)_p-D

wherein: R₁₅ and R₁₆ are individually selected from the group consisting of H, CH₃, C₂-C₁₀ alkyls, C₂-C₁₀ alkenyls or C₂-C₁₀ alkynyls, each of which can be substituted or unsubstituted; straight or branched; and C₂-C₁₀ heteroalkyls, C₂-C₁₀ heteroalkenyls or C₂-C₁₀ heteroalkynyls;

p is a positive integer from 1 to about 12;

D is selected from among -SH, -OH, X₂, -CN, -OR₁₉, NHR₂₀,



wherein:

5

R₁₇ is H, a CH₃ or X₃;

R₁₈ is H, a C₁₋₄ alkyl or benzyl;

R₁₉ is H, a C₁₋₄ alkyl, X₂ or benzyl;

R₂₀ is H, a C₁₋₁₀ alkyl or -C(O)R₂₁

wherein R₂₁ is H, a C₁₋₄ alkyl or alkoxy, t-butoxy or

10

benzyloxy;

X₂ and X₃ are independently selected halogens;

R₃ is H, CH₃, or -C(=O)(CR₁₅R₁₆)_w-D,

where w is 0 or an integer from 1 to about 12, and D is H or as described for R₁ and R₂

J is O, NH or S;

15

R₄, R₅ and R₆ independently selected from the group consisting of H, CH₃, C₂-C₁₀ alkyls, C₂-C₁₀ alkenyls or C₂-C₁₀ alkynyls, each of which can be substituted or unsubstituted; straight or branched; C₂-C₁₀ heteroalkyls, heteroalkenyls or heteroalkynyls and halogens;

R₇ is selected from among H, CH₃ and C₂-C₁₀ alkyls;

20

X₁ is O, NH, or S; and

A is H or a second active moiety;

with a compound of the Formula (V):

(V) L₁-B₁

25

wherein L₁ is a moiety containing a functional group capable of reacting with the NHR₂₂ of Formula (IV);

and B₁ is selected from the group consisting of polymers, biologically active materials and polymeric supports.

1/6

FIG. 1




FIG. 2





FIG. 3








FIG. 4

FIG. 5



6/6

FIG. 6



**(19) World Intellectual Property Organization
International Bureau**



(43) International Publication Date
24 October 2002 (24.10.2002)

PCT

(10) International Publication Number
WO 02/083030 A3


- (1) **International Patent Classification⁷:** C12N 13/00, A01N 51/00, 43/58, 47/10, C07D 241/04, C07C 321/00, 211/00, 49/76
- (21) **International Application Number:** PCT/US02/11476
- (22) **International Filing Date:** 12 April 2002 (12.04.2002)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
60/284,308 17 April 2001 (17.04.2001) US
10/066,323 31 January 2002 (31.01.2002) US
- (71) **Applicant:** CRYOLIFE, INC. [US/US]; 1655 Roberts Boulevard, N.W., Kennesaw, GA 30144 (US).
- (72) **Inventors:** GILBERT, Carl, W.; 4655 Oakleigh Manor Drive, Powder Springs, GA 30127 (US). MCGOWAN, Eleanor, B.; 3062-I Spring Hill Parkway, Smyrna, GA 30080-4735 (US). BLACK, Kirby, S.; 1371 Peppergrass Trail, Acworth, GA 30101 (US). HARPER, T., Gregory, P.; 86 West Euclid Avenue, Stockton, CA 95204 (US).
- (74) **Agent:** MERCANTI, Michael, N.; Roberts & Mercanti, L.L.P., P.O. Box 3156, 105 Lock Street, Suite 203, Newark, NJ 07103-0156 (US).
- (81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
- (88) **Date of publication of the international search report:**
18 December 2003

Published:
— *with international search report*


(88) Date of publication of the international search report:
18 December 2003

[Continued on next page]

(54) Title: BIFUNCTIONAL ENERGY-REVERSIBLE ACYL-COMPOSITIONS



(A)



(B)

(57) Abstract: Energy-reversible acyl conjugates, intermediates, and related compositions are disclosed. In preferred aspects, examples of such compositions include Formulac (A) and (B).

WO 02/083030 A3

WO 02/083030 A3



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/11476

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12N 13/00; A01N 51/00, 43/58, 47/10; C07D 241/04; C07C 321/00, 211/00, 49/76

US CL : 435/173; 514/157, 247, 477; 544/386, 391; 564/161, 162, 305; 568/334

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/173; 514/157, 247, 477; 544/386, 391; 564/161, 162, 305; 568/334

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,114,851 A (PORTER et al) 19 May 1992, entire document	1-31
A	US 5,218,137 A (PORTER et al) 8 June 1993, entire document	1-31
A	US 5,266,453 A (MATUSHITA et al) 30 November 1993, entire document	1-30

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

10 March 2003 (10.03.2003)

Date of mailing of the international search report

10 APR 2003

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Sikari Witherspoon

Telephone No. 703-308-1235

INTERNATIONAL SEARCH REPORT

PCT/US02/11476

Continuation of B. FIELDS SEARCHED Item 3:

REGISTRY, CAPLUS, BIOSIS

search terms: acyl enzymes, energy, light, amino acid, protease, treat, pharmaceutical, cinnamate